

*Christine Nattrass (Yale University)
for the STAR Collaboration*



Outline

- Introduction
- The *Jet* – energy, system, and particle type dependence
- The *Ridge* – energy, system, and particle type dependence
- Comparison to models
- Conclusion


Introduction

Di-hadron correlations


- Define a high- p_T trigger particle
- Look at the distribution of particles relative to that trigger particle



Motivation – Jet and Ridge



- Long-range pseudorapidity ($\Delta\eta$) correlations observed by STAR in Au+Au at intermediate p_T
- Near side jet peak sits on plateau (*Ridge*)
- Significant contribution to the near-side yield in central Au+Au



Method: Yield extraction

- Ridge previously observed to be independent in $\Delta\eta$ in Au+Au
- To determine relative contributions, find yields for near-side, take $\Delta\Phi$ projections in

- $-0.75 < \Delta\eta < 0.75$ **Jet + Ridge**

- $0.75 < |\Delta\eta| < 1.75$ **Ridge**

- $Jet = (Jet+Ridge) - Ridge * .75/1.0$

- $Ridge = \text{yield from } -1.75 < \Delta\eta < 1.75 - Jet \text{ yield}$

- Flow contributions to Jet cancel



- v_2 independent of η for $|\eta| < 1$

- Phys. Rev. C72, 051901(R) (2005), Phys. Rev. Lett. 94, 122


- Yield: number of particles in Jet , $Ridge$

Au+Au 0-10% STAR preliminary


nucl-ex/0701074



The Jet





Jet is like p+p




- Spectra of particles associated with *Jet* harder than inclusive

Jet composition





M. Lamont (STAR), J.Phys.G32:S105-S114,2006
 J. Bielcikova (STAR), v:0707.3100 [nucl-ex]
 C. Nattrass (STAR), arXiv:0804.4683/nucl-ex





- Baryon/meson ratios in *Jet* in Cu+Cu and Au+Au similar to p+p for both strange and non-strange particles

p_T^{trigger} dependence



- No system dependence observed in the data
- Pythia 8.1 describes trends in data up to a scaling factor
 - Gets energy dependence right
 - Stronger deviations at low p_T^{trigger} , as expected

Pythia comparisons






- What can Pythia tell us?

- Higher z_T (lower jet energy) in 62 GeV for same p_T^{trigger}

$p_{T\text{Hat}}$ = the parameter in Pythia for the transverse momentum in the hard subprocess

Pythia comparisons





- What can Pythia tell us?

- Higher z_T (lower jet energy) in 62 GeV for same p_T^{trigger}

p_{THat} = the parameter in Pythia for the transverse momentum in the hard subprocess

p_T associated dependence





- No system dependence
- Pythia 8.1 slightly harder than data
- Diverges slightly from Pythia 8.1 at lower $p_T^{\text{associated}}$

Inverse slope parameter

	$\sqrt{s_{\text{NN}}} = 62$ GeV	$\sqrt{s_{\text{NN}}} = 200$ GeV
Cu+Cu	317 ± 26	445 ± 20
Au+Au	355 ± 21	478 ± 8
d+Au		469 ± 8
Pythia	417 ± 9	491 ± 3

Statistical errors only


N_{part} dependence



- Cu+Cu 62 GeV
- Au+Au 62 GeV
- ★ d+Au 200 GeV
- Cu+Cu 200 GeV
- Au+Au 200 GeV
- Pythia 62 GeV*2/3
- Pythia 200 GeV*2/3


- No system dependence
- Some deviations from Pythia 8.1 with increase in N_{part}
 - Incomplete *Ridge* subtraction?
 - *Jet* modification at low p_T ?

Conclusions: Jet




- Particle ratios similar to p+p
 - Separation of *Jet* and *Ridge* works
 - Effects of triggers which don't come from jets small
 - Pythia can be used to estimate z_T distributions, jet energy
 - Pythia describes data well
 - Scaling factor needed but Pythia 8.1 is not as tuned as earlier versions
 - Energy dependence in *Jet* is described well in Pythia
 - Trends for p_T^{trigger} , p_T^{assoc} dependence right
- *Jet* production mechanism dominated by fragmentation

The Ridge





Ridge is like bulk




- Spectra of particles associated with *Ridge* similar to inclusive

Ridge composition




M. Lamont (STAR), J.Phys.G32:S105-S114,2006
 J. Bielcikova (STAR), v:0707.3100 [nucl-ex]
 C. Nattrass (STAR), arXiv:0804.4683/nucl-ex




- Baryon/meson ratios in *Ridge* similar to bulk for both strange and non-strange particles

Identified trigger: Near-side Yield vs N_{part}




$3.0 \text{ GeV}/c < p_T^{\text{trigger}} < 6.0 \text{ GeV}/c$; $1.5 \text{ GeV}/c < p_T^{\text{associated}} < p$

Ridge yield -
 No trigger type
 dependence



Au+Au $\sqrt{s_{NN}}=200 \text{ GeV}$ from nucl-ex/0701047
 Cu+Cu $\sqrt{s_{NN}}=200 \text{ GeV}$ from SQM2007
 Data points at same N_{part} offset for visibility



Ridge yield vs. $p_{t, trig}$ in Au+Au



Putschke
WWND08



- Ridge yield persists to highest trigger $p_t \Rightarrow$ correlated with jet production

Ridge vs N_{part}



- No system dependence at given N_{part}

Ridge vs N_{part}




- No system dependence at given N_{part}
- *Ridge/Jet* Ratio independent of collision energy



Jet/Ridge w.r.t. reaction plane

Feng QM08

Konzer QM09



20-60%
Au+Au
200 GeV




- Ridge yield decreases with Ψ_s . Smaller ridge yield at larger Ψ_s
- Jet yield approx. independent of Ψ_s and comparable with d+Au

Jet yield independent of Ψ_s , consistent with vacuum fragmentation after energy loss and lost energy deposited in ridge, if medium is “black” out-of-plane and more “gray” in-plane for surviving jets.


Ridge asymmetric in $\Delta\phi$, consistent with surface emission

3-particle correlations




$$\Delta\eta_1 = A1-T$$

$$\Delta\eta_2 = A2-T$$




0-12% d+Au



Netrakanti QM09

$$3 < p_T^{\text{trigger}} < 10 \quad 1 < p_T^{\text{assoc}} < 3 \quad |\Delta\phi| < 0.7$$


- Ridge appears uniform event-by-event within STAR acceptance
- Charge sign dependence:
 - All signs the same: no Jet, Ridge only
 - Friday Plenary session talk (Netrakanti)



Long flow picture bias

S. Vafek et al. / PRL 93 (2004) Phys. A749, 287

Conclusions: Ridge



- Extensive data on Ridge
 - Cu+Cu, Au+Au consistent at same N_{part}
 - *Ridge/Jet* ratio independent of energy
 - Persists to high $p_T^{trigger}$
 - *Ridge* looks like bulk
 - $p_T^{associated}$ dependence, particle composition
 - Appears isotropic in $\Delta\eta$

Comparisons to models

Models


- Radial flow+trigger bias

S. Voloshin, nucl-th/0312065, Nucl. Phys. A749, 287

C.. Pruneau, S. Gavin, S. Voloshin, arXiv:0711.1991v2

E. Shuryak, *Phys.Rev.C*76:047901,2007

- Works for one set of kinematic cuts in central Au+Au at 200 GeV
- Need more detailed comparisons (energy dependence)
- Model needs some refinements (momentum conservation)




- Plasma instability

QCD magnetic fields, Majumder et al, *Phys.Rev.Lett.*99:042301,2007

Anisotropic plasma, P. Romatschke, *PRC*,75014901 (2007)

- So far unable to make enough *Ridge* without Radial flow+trigger bias



Models

- Longitudinal flow

Longitudinal flow, Armesto et al, PRL 93 (2004)

- Problems due to $\Delta\eta$ width

- Momentum kick



Momentum kick from jet, C.-Y. Wong, Phys.Rev.C76:054908,2007

- Fits data well, including energy dependence

- Recombination

Medium heating + recombination, Chiu & Hwa, PRC72, 034903

- No quantitative comparisons



Conclusions

- Considerable evidence that *Jet* is dominantly produced by fragmentation
 - Can we use this information to learn more about the *Ridge*?
- Several models for the *Ridge*, few quantitative comparisons
 - Several depend on hydrodynamics
 - Need better calculations – more quantitative, more than central Au+Au
- Future:
 - More energy dependence (RHIC beam energy scan, LHC)
 - Jet reconstruction – more detailed studies of *Ridge*?
 - Ridge in γ -jet correlations?