

---

---

# *Studies of Jets through Correlations in Heavy Ion Collisions at RHIC*

*Christine Nattrass  
Yale University*


# *Introduction*

---

- Why study jets in heavy ion collisions?
- The near-side
- The away-side
- Conclusions


# Why study jets in heavy ion collisions?

- Hard parton scattering  $\Rightarrow$  back-to-back jets
  - Good (calibrated?) probe of the medium
- High multiplicity in A+A collisions
  - Individual jets cannot be reconstructed
  - Study jets via correlations of particles in space




space

- both azimuth and pseudorapidity




# Why study jets in heavy ion collisions?

- Hard parton scattering  $\Rightarrow$  back-to-back jets
  - Good (calibrated?) probe of the medium
- High multiplicity in A+A collisions
  - Individual jets cannot be reconstructed
  - Study jets via correlations of particles in space



in space

- both azimuth and pseudorapidity





# Why study jets in heavy ion collisions?

- Hard parton scattering  $\Rightarrow$  back-to-back jets
  - Good (calibrated?) probe of the medium
- High multiplicity in A+A collisions
  - Individual jets cannot be reconstructed
  - Study jets via correlations of particles in space



space

- both azimuth and pseudorapidity



# Motivation



- Initial studies showed suppression of away-side peak in A+A collisions
  - $2.0 \text{ GeV}/c < p_T^{\text{associated}}$
- Inclusive  $p_T^{\text{associated}}$ 
  - reappearance of away-side
  - more complex structure than d+Au, p+p




# RHIC

- Luminosity  $\sim 2 \cdot 10^6 \text{ cm}^{-2} \text{ s}^{-1}$
- Can collide multiple species (Au+Au, Cu+Cu, d+Au, p+p...)
- Interaction diamond  $< 20 \text{ cm}$  long
- 56 bunches
- $10^9$  ions/bunch
- nominal 10 hour store time






- Primary Detectors: Drift chambers, time expansion chambers, pad chambers, vertex detector
- Azimuthal coverage: Two sections of  $0 < \varphi < \pi/2$
- Pseudorapidity coverage:  $-0.35 < \eta < 0.35$




- Primary Detector: TPC
- Full azimuthal coverage:  $0 < \varphi < 2\pi$
- Pseudorapidity coverage:  $-1 < \eta < 1$

# Near-side: Motivation

- Near-side shows modification
- Excess yield in Au+Au relative to p+p



# Near-side: Motivation




No  $v_2$  subtraction – signal visible above  $v_2$

- Long-range pseudorapidity ( $\Delta\eta$ ) correlations observed by STAR in Au+Au at intermediate  $p_T$
- Near side jet peak sits on plateau (*Ridge*)
- Significant contribution to the near-side yield in central Au+Au

# Near-side: Ridge production mechanisms

- Parton radiates energy before fragmenting and couples to the longitudinal flow

- gluon bremsstrahlung of hard-scattered parton
- parton shifted to lower  $p_T$
- radiated gluon contributes to broadening





Armesto et al, PRL 93 (2004), nucl-ex/0405301

- Medium heating + parton recombination (Chiu & Hwa, Phys. Rev. C72:034903,2005)

- Recombination of thermal partons only indirectly affected by hard scattering, not part of the jet


- Radial flow + trigger bias (Voloshin nucl-th/0312065, S. A. Voloshin, Nucl. Phys. A749,287 (2005))

# *pT* trigger dependence of Jet and Ridge




- *Jet* yield per trigger increases with  $p_T^{\text{trigger}}$
- Expected because higher  $p_T^{\text{trigger}}$  should be “jettier”
- *Ridge* yield decreases
- *Ridge* dominant in central Au+Au


# Near-side: particle dependence



J. Bielcikova, WWND07



- Identified particles show similar yield trends to unidentified
- Particle ratios in *Ridge* closer to bulk than those in the *Jet*



# Near-side Yield vs $N_{part}$ Cu+Cu vs Au+Au

Identified triggers:


- *Jet* yield

- Nearly flat with  $N_{part}$  within errors across d+Au, Cu+Cu, Au+Au
- No  $v_2$  or background error due to method
- No trigger dependence within errors

- *Ridge* yield

- No *Ridge* within errors in d+Au
- Rises with  $N_{part}$  in Cu+Cu and Au+Au
- No trigger dependence within errors

3.0 GeV/c <  $p_T^{trigger}$  < 6.0 GeV/c; 1.5 GeV/c <  $p_T^{associated}$  <  $p_T^{trigger}$   
C. Nattrass SQM07



Data points at same  $N_{part}$  offset for visibility  
Jet yields: 10% error added to  $V^0$  and h triggers to account for track merging, 15% to  $\Xi$  triggers

# Near-side Yield vs $N_{part}$ Cu+Cu vs Au+Au

Identified triggers:


- *Jet* yield

- Nearly flat with  $N_{part}$  within errors across d+Au, Cu+Cu, Au+Au
- No  $v_2$  or background error due to method
- No trigger dependence within errors

- *Ridge* yield


- No *Ridge* within errors in d+Au
- Rises with  $N_{part}$  in Cu+Cu and Au+Au
- No trigger dependence within errors

3.0 GeV/c <  $p_T^{trigger}$  < 6.0 GeV/c; 1.5 GeV/c <  $p_T^{associated}$  <  $p_T^{trigger}$   
C. Nattrass SQM07



Data points at same  $N_{part}$  offset for visibility  
Jet yields: 10% error added to  $V^0$  and h triggers to account for track merging, 15% to  $\Xi$  triggers


# Near-side: Ridge



- Spectra of particles in *Jet* harder than those of particles in the *Ridge*
- Particles in *Ridge* similar to bulk


# Near-side: Summary

- Enhanced yield in Au+Au collisions relative to p+p
- Extra yield is in *Ridge*
- Particles in Ridge closer to those in the bulk than those in the Jet
  - Particle ratios
  - Spectra
- Consistent with  $N_{\text{part}}$  dependence on *Ridge* size
- Predictions which might distinguish mechanisms not quantitative enough to make detailed comparisons to theory





# Away-side

- Shape change on away-side
- Excess yield at low  $p_T$  on away-side




# Motivation: Away-side



- STAR and PHENIX: qualitative agreement
- STAR: dependent on systematic errors
  - systematic errors from disagreement of different methods of measuring  $v_2$
- PHENIX: claim smaller systematic errors
  - systematic errors from one  $v_2$  measurement (reaction plane)


# Away-side: Motivation

Or STAR results...



## Centrality


- Shape distortion increases with centrality
- Decreases with increasing  $p_T^{\text{trigger}}$
- Degree of shape distortion dependent on  $v_2$  subtraction



- Also see “punch through” of away-side at high  $p_T^{\text{associated}}$


# Away-side: Motivation

Or STAR results...




Centrality →

- Shape distortion increases with centrality
- Decreases with increasing  $p_T^{\text{trigger}}$
- Degree of shape distortion dependent on  $v_2$  subtraction
- Also see “punch through” of away-side at high  $p_T^{\text{associated}}$



# Away-side: Motivation

Or STAR results...




Centrality →

- Shape distortion increases with centrality
  - Decreases with increasing  $p_T^{\text{trigger}}$
  - Degree of shape distortion dependent on  $v_2$  subtraction
  - Also see “punch through” of away-side at high  $p_T^{\text{associated}}$
- AuAu  
● AuAu  $|\Delta\eta| < 0.72$   
● dAu

# Away-side: Motivation

Or STAR results...



Centrality →

- Shape distortion increases with centrality
  - Decreases with increasing  $p_T^{\text{trigger}}$
  - Degree of shape distortion dependent on  $v_2$  subtraction
  - Also see “punch through” of away-side at high  $p_T^{\text{associated}}$
- AuAu  
● AuAu  $|\Delta\eta| < 0.72$   
● dAu

# Away-side: Proposed models

---


- Where could this come from?
  - Large angle gluon radiation (Vitev and Polsa and Salgado).
  - Deflected jets, due to flow (Armesto, Salgado and Wiedemann) and/or path length dependent energy loss (Chiu and Hwa).
  - Hydrodynamic conical flow from mach cone shock-waves (Stöcker, Casalderrey-Solanda, Shuryak and Teaney, Renk, Ruppert and Muller).
  - Cerenkov gluon radiation (Dremin, Koch).
- 2-particle correlations give the same qualitative results for all models → can't distinguish

# 3-particle correlations


- Two different coordinate systems
- Different background subtractions
- Both assume 2 components (jet + flow background)

STAR

$\Delta\phi$ - $\Delta\phi$  space ( $\Delta\phi = \phi - \phi_{\text{Trigger}}$ )




- PHENIX
- polar coordinates



Ajitanand (PHENIX) HP06, IWCF'06

# Conclusions

- Jets have proven to be a powerful probe at RHIC
  - The near-side is modified
    - The Jet
      - Looks like vacuum fragmentation
    - The Ridge
      - Looks very similar to the bulk
  - Away-side is modified
- Many proposed models, more work to do



# Wisdom from Goethe

---

Delacroix



Da steh' ich nun, ich armer Tor,  
Und bin so klug als wie zuvor!


Faust I

Goethe



# Away-side: Motivation

PHENIX  
Au+Au & p+p  
 $\sqrt{s} = 200$  GeV  
arXiv:0705:3238

Whether we  
look at  
PHENIX  
results...





# 3-particle correlations: PHENIX



Need to work on this slide,  
colors, etc.



# 3-particle correlations: PHENIX



Near-side

Need to work on this slide,  
colors, etc.




# 3-particle correlations: PHENIX




Need to work on this slide,  
colors, etc.

Away-side


# 3-particle correlations: PHENIX







Normal jet simulations



Deflected jet simulations




Mach Cone simulations





# 3-particle correlations: PHENIX

Ajitanand (PHENIX) HP06, IWCF'06









## $\Delta\phi^*$ Projections

$v_2$  subtracted




- Dependent on background subtraction
- Details of background subtraction not shown
- Central data not shown

# 3-particle correlations: STAR



# 3-particle correlations: STAR






Subtracting a complicated background...

Still assuming ZYAM...



$$v_4 = 1.15 v_2^2$$



# 3-particle Correlations: STAR Results






J. Ulery, ISMD07



Structure changes with centrality


Peripheral Au+Au looks like d+Au, p+p

Central Au+Au shows new features


 Near-Near  
  Near-Away  
  Away-Away  
 $1 < p_{\text{T}}^{\text{associated}} < 2 \text{ GeV}/c, 3 < p_{\text{T}}^{\text{trigger}} < 4 \text{ GeV}/c$

# 3-particle Correlations: STAR Results

- On-diagonal and off-diagonal projections.
- Yellow bands are systematic errors.
- Significant off-diagonal peaks.







# 3-particle Correlations: STAR Results



- Attempts at extracting emission angle
- No apparent  $p_T$  dependence
- No apparent centrality dependence

# 3-particle Correlations: *STAR* Cumulant




- Clear mathematical definition
- Difficult to interpret
- Favors modification but the type of modification is unclear

# *3-particle Correlations: Summary*


---

- Both STAR and PHENIX standard analyses slightly favor conical emission
  - STAR data slightly favors Mach cone over Cerenkov radiation
  - Both analyses dependent on validity of
    - ZYAM
    - 2 component picture
- Similar STAR analysis which does not support conical emission but has unclear interpretation

# Another method - Baryon/Meson Ratio




- If there is a Mach cone, angle should depend on mass
- Error bars are still too large to conclude
- However, at least systematic errors for  $K_S^0$  and  $\Lambda$  are correlated




- Disadvantages:
  - Current error bars prevent any conclusions
- Advantages:
  - STAR year 7 data allow 4x statistics
  - New method (Brooke Haag, QM08 poster) may allow even higher stats/higher  $p_T$
  - systematic errors due to  $v_2$  at least move together


# Summary






- Studies showing shape changes on away-side
  - Signal/Background  $> 1/20$
  - Systematic errors due to  $v_2$  large because background is large
  - $1 \text{ GeV/c} < p_T^{\text{associated}}$ 
    - Ridge yield significant on Near-side
    - $\rightarrow$  ZYAM assumption could lead to significant errors



- We don't understand the near-side yet
  - 2.5 GeV/c trigger (PHENIX) isn't very "jetty"
  - 3.0 GeV/c (STAR) is only slightly better
  - $1 \text{ GeV/c} < p_T^{\text{associated}}$  - on near-side, Ridge dominates. Does this affect the away-side?



- How do we know that the production mechanisms for the Ridge and the shape changes on the away-side are distinct?



# Outlook

---

- Higher  $p_T$  on 3-particle correlations?
  - Really need higher  $p_T^{\text{associated}}$ ,  $p_T^{\text{trigger}}$  to understand results
  - More data not yet analyzed
  - Detector upgrades which would allow more data in a run
  - Advances in triggering
- Need to understand the near-side
  - More data helps in particle ratios
  - More particles?
- Jet reconstruction in heavy ion collisions?
  - Some progress
  - Would allow reconstruction of near-side


# *Backup slides*

---



# Reminder: Background Subtraction

Phys.Rev.Lett.98:232302,2007



No dip before background subtraction

Signal sits on top of large background


Assumption that there is no yield at minimum

Dip appears after background subtraction





# A caveat...

- Large background subtraction...
  - Signal/Background  $\approx 0.05$ 
    - Depends on kinematic region
    - Signal/Background higher at higher  $p_T$



# Determination of yields and errors



- Background:

$$B(1 + 2 v_2^{\text{trig}} v_2^{\text{assoc}} \cos(2\Delta\Phi))$$

$v_2$  – elliptic flow

- Different fit methods for determination of B

- Assume there is no yield correlated with the jet at some point

- Zero Yield At Minimum (ZYAM)

- Zero Yield At 1 (ZYA1)

- $v_2$  error -> systematic error on correlations *assuming ZYAM is correct*

# ***Assumptions in background subtraction***


---

- The only background is elliptic flow
- Elliptic flow is independent of jets and therefore the correlations can be separated into two *independent* components
- There is a point in azimuth where none of the correlations are due to jets

**Any conclusions are heavily dependent on the validity of these assumptions!**

# Near-side: Method

- *Ridge* previously observed to be flat in  $\Delta\eta$  in Au+Au
- To determine relative contributions, find yields for near-side, take  $\Delta\Phi$  projections in
  - $-0.75 < \Delta\eta < 0.75$  *Jet + Ridge*
  - $0.75 < |\Delta\eta| < 1.75$  *Ridge*
  - $Jet = (Jet+Ridge) - Ridge * .75/1.75$
  - $Ridge = \text{yield from } -1.75 < \Delta\eta < 1.75 - Jet \text{ yield}$
- Flow contributions to jet cancel
- $v_2$  flat with  $\eta$  for  $|\eta| < 1$




• Phys. Rev. C72, 051901(R) (2005), Phys. Rev. Lett. 94, 122303 (2005)

# $p_T$ -distribution of associated particles

- Ridge spectra similar to the bulk
  - Cu+Cu measurements probably not possible
- Jet spectra are slightly harder
  - Cu+Cu T within error of Au+Au

Trigger particle	T (ridge) MeV	T (jet) MeV
$h^{+/-}$	$438 \pm 4$ (stat.)	$478 \pm 8$
$K^0_s$	$406 \pm 20$ (stat.)	$530 \pm 61$
$\Lambda$	$416 \pm 11$ (stat.)	$445 \pm 49$

nucl-ex/0701047



Fit to  $A \exp(-p_T/T)$

# Two Analysis Techniques

Measure 1-, 2-, and 3-Particle Densities

$$\rho_1(j_i) \propto \frac{d^2 N}{dj_i} \quad \rho_2(Dj_{ij}) \propto \frac{d^2 N}{dDj_{ij}} \quad \rho_3(Dj_{ij}, Dj_{ik}) \propto \frac{d^3 N}{dDj_{ij} dDj_{ik}}$$

3-particle densities = superpositions of **truly** correlated 3-particles, and combinatorial components.

We use **two approaches** to extract the truly correlated 3-particles component

- Cumulant technique:

$$C_3(\Delta\phi_{12}, \Delta\phi_{13}) = \rho_3(\Delta\phi_{12}, \Delta\phi_{13}) - \rho_2(\Delta\phi_{12})\rho_1(3) - \rho_2(\Delta\phi_{13})\rho_1(2) - \rho_2(\Delta\phi_{13} - \Delta\phi_{12})\rho_1(1) + 2\rho_1(1)\rho_1(2)\rho_1(3)$$

**PROs** Simple Definition  
Model Independent.

**CONS** Not positive definite  
Interpretation perhaps difficult.


- Jet+Flow Subtraction Model:

$$\bar{J}_3(\Delta\phi_{12}, \Delta\phi_{13}) = v_3(\Delta\phi_{12}, \Delta\phi_{13}) - \bar{v}_2^*(\Delta\phi_{12})B_2(\Delta\phi_{13}) - \bar{v}_2^*(\Delta\phi_{13})B_2(\Delta\phi_{12}) - B_3(\Delta\phi_{12}, \Delta\phi_{13})$$

Intuitive in concept  
Simple interpretation in principle.


Model Dependent  
 $v_2$  and normalization factors  
systematics


# Measurement of 3-Particle Cumulant



- Clear evidence for **finite 3-Part Correlations**
- Observation of flow like and jet like structures.
  - Evidence for  $v_2 v_2 v_4$  contributions

# 3-Cumulant vs. centrality






# 3-particle correlations: STAR


Ulery (STAR) QM05

- Trigger particle  $3 < p_T < 4$  GeV/c with pairs of associated particle  $1 < p_T < 2$  GeV/c..
- Complicated background...
  - Raw signal contains (Jet+Bkgd)  $\otimes$  (Jet+Bkgd).
  - To obtain Jet  $\otimes$  Jet we must subtract Bkgd  $\otimes$  Bkgd and Jet  $\otimes$  Bkgd (and Bkgd  $\otimes$  Jet.)



# Jet and Jet+Ridge yields & widths


Correlate Jet ( $\Delta\eta(J)$ ) and Jet+Ridge ( $\Delta\phi(J+R)$ ) widths & yields via centrality



- Jet+Ridge yield increasing with centrality
- Jet+Ridge shape asymmetric in  $\Delta\eta$  and  $\Delta\phi$

# Jet yields & widths: $\Delta\eta$ vs. $\Delta\phi$


Correlate Jet ( $\Delta\eta(J)$ ) and Jet ( $\Delta\phi(J)$ ) widths and yields via centrality



- Jet yield  $\sim$  symmetric in  $\Delta\eta \times \Delta\phi$
- Jet shape  $\sim$  symmetric in  $\Delta\eta \times \Delta\phi$  for  $p_{t, \text{trig}} > 4$  GeV  
(asymmetric in  $\Delta\eta$  for  $p_{t, \text{trig}} < 4$  GeV)

# A caveat...

- Large background subtraction




# ***3-particle correlations: STAR background***

---

- Add page(s) w/ equations

# Jet and Jet+Ridge yields & widths


Correlate Jet ( $\Delta\eta(J)$ ) and Jet+Ridge ( $\Delta\phi(J+R)$ ) widths & yields via centrality



- Jet+Ridge yield increasing with centrality
- Jet+Ridge shape asymmetric in  $\Delta\eta$  and  $\Delta\phi$

# Jet yields & widths: $\Delta\eta$ vs. $\Delta\phi$

Correlate Jet ( $\Delta\eta(J)$ ) and Jet ( $\Delta\phi(J)$ ) widths and yields via centrality



- Jet yield  $\sim$  symmetric in  $\Delta\eta \times \Delta\phi$
- Jet shape  $\sim$  symmetric in  $\Delta\eta \times \Delta\phi$  for  $p_{t,\text{trig}} > 4$  GeV  
(asymmetric in  $\Delta\eta$  for  $p_{t,\text{trig}} < 4$  GeV)

# Away-side: Motivation

---

- See shape change in 2-D correlations
- apparent peak at  $d\phi \neq \pi$
- already see strong dependence on the background subtraction
- Qualitatively STAR and PHENIX see the same thing
- shape change as you go peripheral  $\rightarrow$  central
- quantitatively STAR and PHENIX disagree dramatically on size of systematic errors

# *3-particle correlations*

---

- Two different approaches to measurement w/ different ways of plotting data
- Similar qualitative results – weak support for conical emission
- Statistical errors - can be reduced
- Systematic errors – can they be reduced?